

## I. AMENDMENTS TO THE SPECIFICATION

Please replace paragraphs as filed with the following replacement paragraphs:

**[00154]** Examples of cleavable oligonucleotides which contain two reverse U (rU) linkers and have been synthesized on a chip are as follows:

Probe	Pu1	PS1	PU2	PS2
IL6_T7	5'CAAGGATCTTACCGCTGTTGtgaggagacttgcctggtrUTAATACGACTCACTAT AGGt ctgcaggactggatcaggrU ( <u>SEQ ID NO:1</u> )			
CYP11A_t7	5'CAAGGATCTTACCGCTGTTGgtgaccctgcagagatatctrUTAATACGACTCACTA TAGGtgtccggaa taggtgatgtrU( <u>SEQ ID NO:2</u> )			
ATP2A1_T7	5'CAAGGATCTTACCGCTGTTGgattggcattgccatggatrUTAATACGACTCACTAT AGGtccacagcagctacgatggrU ( <u>SEQ ID NO:3</u> )			
IL6_Nick	5'caaggattt accgctgttg tgaggagact tgcctggtn cgctccagac ttgagtccga tctgcaggaa ctggatcaggrU ( <u>SEQ ID NO:4</u> )			
CYP11A_Nick	5'CAAGGATCTTACCGCTGTTGgtgaccctgcagagatatctrUCGCTCCAGACTTGAG TCCGAgttccggaa gtaggtgatgtrU ( <u>SEQ ID NO:5</u> )			
ATP2A1_Nick	5'CAAGGATCTTACCGCTGTTGgattggcattgccatggatrUCGCTCCAGACTTGAG TCCGAtccacagcagctacgatggrU ( <u>SEQ ID NO:6</u> )			

**[00164]** Different strategies can be used to release or cleave oligonucleotides synthesized on a solid substrate from that substrate. The cleavage efficiency of three different linkers was examined to determine the preferred linker(s) for cleaving oligonucleotides from a solid substrate (rU is 5'-phosphoramidite with 2'-acetyl and 3'DMT; U is 3'-phosphoramidite with 2'fpmp and 5'-DMT; and dU is 2'-deoxyuridine). To begin, the following oligonucleotides were synthesized using an Expetide™ DNA synthesizer and standard phosphoamidite chemistry:

Sequence A      3' tttttttttrugtccacagcatccga-fam-5' (SEQ ID. NO:7)

Sequence B      3' tttttttttugtccacagcatccga-fam-5' (SEQ ID. NO:8)

Sequence C      3' tttttttttdugtccacagcatccga-fam-5' (SEQ ID. NO:9)

[00168] The GFP gene is 714 base pairs (bp) long. Suitable subchains (computational fragmentation) for the assembly of the GFP gene were selected, and oligonucleotides between 40 and 47 nucleotides long were synthesized on a chip using the methods outlined above. The complete set of 34 GFP subchains synthesized on a chip are as follows:

GFP-F2	ATGAGTAAAG GAGAAGAACT TTCACTGGA GTTGTCCCAA TTCTTG	<u>SEQ ID NO:10</u>
GFP-F3	TTGAATTAGA TGGTGTATGTT AATGGGCACA AATTTCTGT CAGT	<u>SEQ ID NO:11</u>
GFP-F4	GGAGAGGGTG AAGGTGATGC AACATACGGA AAACCTACCC T	<u>SEQ ID NO:12</u>
GFP-F5	TAAATTATT TGCACTACTG GAAAACCTACC TGTCCATGG CCAA	<u>SEQ ID NO:13</u>
GFP-F6	CACTTGTACAC TACTTCTCT TATGGTGTTC AATGCTTTTC AAGATA	<u>SEQ ID NO:14</u>
GFP-F7	CCCAGATCAT ATGAAACGCC ATGACTTTT CAAGAGTGCC AT	<u>SEQ ID NO:15</u>
GFP-F8	GCCCCGAAGGT TATGTACAGG AAAGAACTAT ATTTTCAAA GATG	<u>SEQ ID NO:16</u>
GFP-F9	ACGGGAACTA CAAGACACGT GCTGAAGTCA AGTTTGAAGG T	<u>SEQ ID NO:17</u>
GFP-F10	GATACCCCTTG TTAATAGAAC CGAGTTAAAA GGTATTGATT TTAAAG	<u>SEQ ID NO:18</u>
GFP-F11	AAGATGGAAA CATTCTTGGG CACAAATTGG AATACAACTA TAACTC	<u>SEQ ID NO:19</u>
GFP-F12	ACACAATGTA TACATCATGG CAGACAAACA AAAGAATGGA ATCAA	<u>SEQ ID NO:20</u>
GFP-F13	AGTTAACTTC AAAATTAGAC ACAACATTGA AGATGGAAGC GTTCA	<u>SEQ ID NO:21</u>
GFP-F14	ACTAGCAGAC CATTATCAAC AAAATACTCC AATTGGCGAT GG	<u>SEQ ID NO:22</u>
GFP-F15	CCCTGTCCTT TTACCAAGACA ACCATTACCT GTCCACACAA T	<u>SEQ ID NO:23</u>
GFP-F16	CTGCCCTTTC GAAAGATCCC AACGAAAAGA GAGACCACAT G	<u>SEQ ID NO:24</u>
GFP-F17	GTCCTTCTTG AGTTTGTAAAC AGCTGCTGGG ATTACACATG GC	<u>SEQ ID NO:25</u>
GFP-F18	ATGGATGAAC TATACAAATA GCATTCGTAG AATTGACTCT ATAGTG	<u>SEQ ID NO:26</u>
GFP-R1	TGAAAAGTTC TTCTCCTTTA CTCAT	<u>SEQ ID NO:27</u>
GFP-R2	ATTAACATCA CCATCTAATT CAACAAAGAAT TGGGACAACCT CCAG	<u>SEQ ID NO:28</u>
GFP-R3	CATCACCTTC ACCCTCTCCA CTGACAGAAA ATTGTGCCCC	<u>SEQ ID NO:29</u>
GFP-R4	TTTCCAGTAG TGCAAATAAA TTAAAGGGTA AGTTTCCGT ATGTTG	<u>SEQ ID NO:30</u>
GFP-R5	ATAAGAGAAA GTAGTGACAA GTCTGGCCA TGGAACAGGT AGT	<u>SEQ ID NO:31</u>
GFP-R6	GCCGTTTCAT ATGATCTGGG TATCTTGAAA AGCATTGAAC ACC	<u>SEQ ID NO:32</u>
GFP-R7	CCTGTACATA ACCCTCGGGC ATGGCACTCT TGAAAAAGTC AT	<u>SEQ ID NO:33</u>
GFP-R8	ACGTGTCTTG TAGTTCCCGT CATCTTGAA AAATATAGTT CTIT	<u>SEQ ID NO:34</u>
GFP-R9	CGATTCTATT ACAAGGGTA TCACCTICAA ACTTGACTTC AGC	<u>SEQ ID NO:35</u>
GFP-R10	TGTCCAAGAA TGTTCACATC TTCTTTAAAA TCAATACCTT TTAACT	<u>SEQ ID NO:36</u>
GFP-R11	TGCCATGATG TATACATTGT GTGAGTTATA GTTGTATTCC AATTG	<u>SEQ ID NO:37</u>
GFP-R12	TTGTGTCTAA TTTGAAGTT AACTTGATT CCATTCTTT GTTTGTC	<u>SEQ ID NO:38</u>
GFP-R13	TTGTTGATAA TGGTCTGCTA GTTGAACGCT TCCATCTTCA ATG	<u>SEQ ID NO:39</u>
GFP-R14	TGTCTGGTAA AAGGACAGGG CCATGCCAA TTGGAGTATT	<u>SEQ ID NO:40</u>
GFP-R15	GGGATCTTTC GAAAGGGCAG ATTGTGTGGA CAGGTAATGG T	<u>SEQ ID NO:41</u>
GFP-R16	CTGTTACAAA CTCAGAACGG ACCATGTGGT CTCTCTTTTC GTT	<u>SEQ ID NO:42</u>
GFP-R17	TGCTATTGT ATAGTTCATC CATGCCATGT GTAATCCCAG CAG	<u>SEQ ID NO:43</u>

[00169] Additionally, the following two control oligonucleotides (Puc2PM- perfect match and Puc2MM- mismatch) were also synthesized on the chip using the methods outlined above:

PUC2PM	CTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTA	SEQ ID NO:44
PUC2MM	CTGGCAGTAGCCACTGGTAACAGGATTAGCAGAGCGAGGTATGTA	SEQ ID NO:45

[00179] One method for releasing or cleaving synthesized oligonucleotides from a solid substrate is an enzymatic approach involving the use of restriction endonuclease (R.E.) enzymes to selectively and specifically cleave desired oligonucleotides from the substrate surface. To test this approach, the Dpn II R.E. enzyme was used to cleave two complementary oligonucleotide DNAs, the first oligo being GFP-F2Part 5'-CACTGGAGTTGTCCCAATTCTTGatcgcc-3' (SEQ ID NO:46) and the second one being DpnIISite 5'-ggccgatcCAA-3' (SEQ ID NO:47). Since II enzyme recognizes and cleaves the sequence 5'-^GATC-3', the isolation of clean oligonucleotides was expected after digestion with the enzyme. Our initial test on the digested oligonucleotides in solution phase was successful. In the experiment, two oligonucleotides were mixed at a molar ration of 1:5 (GFP-F2Part:DpnIISite) and incubated with or without Dpn II enzyme at 37°C. These reactions were analyzed at various time points with CE (capillary electrophoresis, 10% polyacryliamid gel with 7 M urea). As shown in Figure 23, approximately 80% of the longer oligonucleotides were cut by Dpn II in 1 hour. This experiment demonstrates the efficient release of synthesized oligonucleotides from the substrate surface through the use of R.E. enzymes.

[00180] In other embodiments of the present disclosure, an oligonucleotide sequence can be synthesized such that it will anneal itself, thereby forming a duplex oligonucleotide with a hairpin loop. The duplex DNA can then be digested with an enzyme, for example a R.E. enzyme, to form double-stranded DNA that can be ligated to other double-stranded DNA and/or oligonucleotides. To demonstrate the ability of a R.E. enzyme to digest a synthesized oligonucleotide that anneals to itself, the following oligonucleotide sequences with FAM label (DEFINE FAM) were synthesized on a chip with a regular DMT chip surface:

ePM-40	FAM- CTGGCAGCAGCCACTTAACTATCGGGCATTTAACTATCGGATCGGCCTTTGGCGATCGCAT AGTTAAATGCCGATAGTTAAAGTGGCTGCTGCCAG	SEQ ID NO:48
ePM-20	FAM- CTGGCAGCAGCCACTTAACTATCGGGCATTTAACTATCGGATCGGCCTTTGGCGATCGCAT AGTTAAATGCCGATA	SEQ ID NO:49

eMM-40	FAM-CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATCGCAT AGTTACATGCCCATAGTTAAAGTGGCTGCCAG	<u>SEQ ID NO:50</u>
eMM-40-2	FAM-CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATCGCAT AGTTACATGCCCATAGTTAAAGTGGCTGCCAG	<u>SEQ ID NO:51</u>
eMM-20	FAM-CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATCGCAT AGTTACATGCCCATAGTTAAAGTGGCTGCCAG	<u>SEQ ID NO:52</u>
eD-40	FAM-CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATCGCAT AGTTAATGCCCATAGTTAAAGTGGCTGCCAG	<u>SEQ ID NO:53</u>
eD-40-2	FAM-CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATCGCAT AGTTAATGCCCATAGTTAAAGTGGCTGCCAG	<u>SEQ ID NO:54</u>
eD-20	FAM-CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATCGCAT AGTTAATGCCCATAGTTAAAGTGGCTGCCAG	<u>SEQ ID NO:55</u>

[00183] This efficiency of the PGA chemistry utilized in the present disclosure also results in the ability of this chemistry to generate synthetic oligonucleotide sequences that are significantly longer than those that could be synthesized using previously disclosed methods. A programmable light-directed synthesis system was used to synthesize oligomers up to 100 nucleotides in length on a microfluidic array chip. The oligonucleotides synthesized on a chip were as follows:

Pu2P -100	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCATTAACTATCGGCATTAACTATGC	<u>SEQ ID NO:56</u>
Pu2P M-95	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCATTAACTATCGGCATTAAAC	<u>SEQ ID NO:57</u>
Pu2P M-90	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCATTAACTATCGGCAT	<u>SEQ ID NO:58</u>
Pu2P M-85	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCATTAACTATGC	<u>SEQ ID NO:59</u>
Pu2P M-80	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCATTAAAC	<u>SEQ ID NO:60</u>
Pu2P M-75	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCAT	<u>SEQ ID NO:61</u>
Pu2P M-70	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATGC	<u>SEQ ID NO:62</u>
Pu2P M-65	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCAT	<u>SEQ ID NO:63</u>
Pu2P M-60	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATCGGCAT	<u>SEQ ID NO:64</u>
Pu2P M-55	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAACT ATGC	<u>SEQ ID NO:65</u>
Pu2P M-50	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGCATCGCCTTGGCCGATTTAAAC	<u>SEQ ID NO:66</u>
Pu2P M-45	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATCGGCAT	<u>SEQ ID NO:67</u>
Pu2P M-40	CTGGCAGCAGCCACTTTAACTATCGGCATTAACTATGC	<u>SEQ ID NO:68</u>
Pu2P M-35	CTGGCAGCAGCCACTTTAACTATCGGCATTAAAC	<u>SEQ ID NO:69</u>
Pu2P M-30	CTGGCAGCAGCCACTTTAACTATCGGCAT	<u>SEQ ID NO:70</u>
Pu2P M-25	CTGGCAGCAGCCACTTTAACTATGC	<u>SEQ ID NO:71</u>
Pu2P M-20	CTGGCAGCAGCCACTTTAAC	<u>SEQ ID NO:72</u>

Pu2P M-15	CTGGCAGCAGCCACT	<u>SEQ ID NO:73</u>
Pu2M M-100	CTGGCAGTAGCCACTTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGGCATTTAACT ATGCGGCATTAACCTATGCGGCATTTAACTATGC	<u>SEQ ID NO:74</u>
Pu2M M-95	CTGGCAGTAGCCACTTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACT ATGCGGCATTAACCTATGCGGCATTTAAC	<u>SEQ ID NO:75</u>
Pu2M M-90	CTGGCAGTAGCCACTTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACT ATGCGGCATTAACCTATGCGGCAT	<u>SEQ ID NO:76</u>
Pu2M M-85	CTGGCAGTAGCCACTTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACT ATGCGGCATTAACCT ATGC	<u>SEQ ID NO:77</u>
Pu2M M-80	CTGGCAGTAGCCACTTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAAAC TTAACCTATGCGGCATTTAACCTATGCGCATTAAAC	<u>SEQ ID NO:78</u>
Pu2M M-75	CTGGCAGTAGCCACTTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACT ATGCGGCAT	<u>SEQ ID NO:79</u>
Pu2M M-70	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACT ATGC	<u>SEQ ID NO:80</u>
Pu2M M-65	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAAAC	<u>SEQ ID NO:81</u>
Pu2M M-60	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACT ATGCGCATT	<u>SEQ ID NO:82</u>
Pu2M M-55	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTATGC	<u>SEQ ID NO:83</u>
Pu2M M-50	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGCGCATTAAAC	<u>SEQ ID NO:84</u>
Pu2M M-45	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGCGCATT	<u>SEQ ID NO:85</u>
Pu2M M-40	CTGGCAGTAGCCACTTAACTATGCGCATTAACTATGC	<u>SEQ ID NO:86</u>
Pu2M M-35	CTGGCAGTAGCCACTTAACTATGCGCATTAAAC	<u>SEQ ID NO:87</u>
Pu2M M-30	CTGGCAGTAGCCACTTAACTATGCGCATT	<u>SEQ ID NO:88</u>
Pu2M M-25	CTGGCAGTAGCCACTTAACTATGC	<u>SEQ ID NO:89</u>
Pu2M M-20	CTGGCAGTAGCCACTTAAAC	<u>SEQ ID NO:90</u>
Pu2M M-15	CTGGCAGTAGCCACT	<u>SEQ ID NO:91</u>
Puc2D -100	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGCAGCATTAACCTATGCGCATTAACTATGC	<u>SEQ ID NO:92</u>
Puc2D -95	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGCAGCATTAACCTATGCGCATTAAAC	<u>SEQ ID NO:93</u>
Puc2D -90	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGCAGCATTAACCTATGCGCATT	<u>SEQ ID NO:94</u>
Puc2D -85	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGCAGCATTAACCTATGC	<u>SEQ ID NO:95</u>
Puc2D -80	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGCAGCATTAAC	<u>SEQ ID NO:96</u>
Puc2D -75	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGCAGCAT	<u>SEQ ID NO:97</u>
Puc2D -70	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAACTA TGC	<u>SEQ ID NO:98</u>
Puc2D -65	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATTAAAC	<u>SEQ ID NO:99</u>
Puc2D -60	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCATT	<u>SEQ ID NO:100</u>
Puc2D -55	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAACTATGC	<u>SEQ ID NO:101</u>
Puc2D -50	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATTAAAC	<u>SEQ ID NO:102</u>
Puc2D -45	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGCGCATT	<u>SEQ ID NO:103</u>
Puc2D -40	CTGGCAGAGCCACTTAACTATGCGCATTAACTATGC	<u>SEQ ID NO:104</u>

Puc2D -35	CTGGCAGAGCCACTTTAACTATGCGGCATTTAAC	<u>SEQ ID NO:105</u>
Puc2D -30	CTGGCAGAGCCACTTTAACTATGCGGCAT	<u>SEQ ID NO:106</u>
Puc2D -25	CTGGCAGAGCCACTTTAACTATGC	<u>SEQ ID NO:107</u>
Puc2D -20	CTGGCAGAGCCACTTTAAC	<u>SEQ ID NO:108</u>
Puc2D -15	CTGGCAGAGCCACT	<u>SEQ ID NO:109</u>
Stem- 85	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGGCATTTAACTATGCGGCATTTAACT ATGCGGCATTAACATATGC	<u>SEQ ID NO:110</u>
Stem- 80	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGGCATTTAACTATGCGGCATTTAACT ATGCGGCATTAAC	<u>SEQ ID NO:111</u>
Stem- 75	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGGCATTTAACTATGCGGCATTTAACT ATGCGGCAT	<u>SEQ ID NO:112</u>
Stem- 70	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGGCATTTAACTATGCGGCATTTAACT ATGC	<u>SEQ ID NO:113</u>
Stem- 65	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGGCATTTAACTATGCGGCATTTAAC	<u>SEQ ID NO:114</u>
Stem- 60	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACTATGCGGCAT	<u>SEQ ID NO:115</u>
Stem- 55	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACTATGCGGCATTTAACTATGC	<u>SEQ ID NO:116</u>
Stem- 50	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAAAC	<u>SEQ ID NO:117</u>
Stem- 45	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACTATGCGCAT	<u>SEQ ID NO:118</u>
Stem- 40	TTAACTATGCGGCATTTAACTATGCGCATTAACTATGCGCATTAACTATGC	<u>SEQ ID NO:119</u>
Stem- 35	TTAACTATGCGGCATTTAACTATGCGCATTAAAC	<u>SEQ ID NO:120</u>
Stem- 30	TTAACTATGCGGCATTTAACTATGCGCATT	<u>SEQ ID NO:121</u>
Stem- 25	TTAACTATGCGGCATTTAACTATGC	<u>SEQ ID NO:122</u>
Stem- 20	TTAACTATGCGGCATTTAAC	<u>SEQ ID NO:123</u>
Stem- 15	TTAACTATGCGGCAT	<u>SEQ ID NO:124</u>
Stem- 10	TTAACTATGC	<u>SEQ ID NO:125</u>
Stem-5	TTAAC	<u>SEQ ID NO:126</u>

[00184] The oligonucleotides were designed to contain a 15-mer probe (CTGGCAGCAGCCACT) (SEQ ID NO:73) at their 5'-end and connected to variable sizes of non-probe sequence from 0 to 85 nucleotides in length. Additionally, a single base mismatch 15-mer (CTGGCAGTAGCCACT) (SEQ ID NO:91) probe and a single base deletion 14-mer (CTGGCAGAGCCACT) (SEQ ID NO:109) probe were also synthesized on the chip as control sequences. Oligonucleotides from 5 to 100 nucleotides in length were synthesized on the chip, and the two control sequences were arranged side by side in the array for comparison purpose. After the oligomers were synthesized on the array chip, the chip was deprotected with EDA at room temperature for 2 hours and fill with 6xSSPE buffer. The 15 nucleotide target

oligonucleotide labeled with a Cy3 dye was hybridized to the chip in 6xSSPE for 2 hours at room temperature, and the chip was subsequently washed with 0.001xSSPE buffer. As illustrated in Figure 25 and shown in Figure 26, the presence of fluorescence on the chip after the hybridization assay demonstrates that 100-mer oligonucleotides were synthesized on the chip. Additionally, the fluorescence intensity profile indicated a stepwise yield of 98.5% for the synthesis of these long oligonucleotides, which is a significant improvement over known methods for synthesizing oligonucleotides on an array chip. In another experiment, a comparison of the per step yield for oligonucleotides 15 to 100 nucleotides in length on a dual chip demonstrated an even higher stepwise yield of 98.9% and 99.1% (figure 27).

[00187] The disclosed methods for generating pools of oligomers can also be used to generate an RNAi (RNA interference) chip. 252 oligonucleotides were generated on an RNAi chip using the methods previously outlined, with each oligonucleotide synthesized containing a SAP1 sequence (TGCAGTTAGCTCTTCCASAT) (SEQ ID NO:128) at the 3' end, a variable RNAi specific sequence in the middle (22 nucleotides in length), and a T7 promotor sequence CCTATAGTGAGTCGTATTA) (SEQ ID NO:129) at the 5'-end (total length about 60 nucleotides). In order to cleave the oligonucleotides from the chip, reverse-U was incorporated into the 3'-end of all oligonucleotides. Additionally, the same two control oligonucleotides (puc2PM- (SEQ ID NO:44-perfect match) and Puc2MM- (SEQ ID NO:45-mismatch) as disclosed in example 3 were also synthesized on the RNAi chip. The quality of the oligonucleotides synthesized on the RNAi chip was also analyzed by hybridization with Cy3 labeled 15-mer Puc2 target as outlined in Example 3.

**II. ADDITION OF SEQUENCE LISTING**

Please place the attached paper copy of the "Sequence Listing" in the captioned application beginning as a new page after the "Abstract of the Disclosure."

**III. CONCLUSION**

Applicants believe that a full and complete reply has been made to the Request to Comply with Sequence Rules and, as such, the present application is in condition for allowance. Prompt and favorable consideration of this Amendment and Reply is respectfully requested.

Respectfully submitted,



Margaret J. Sampson  
Reg. No. 47,052  
Attorney for Applicant

Vinson & Elkins L.L.P.  
First City Tower  
1001 Fannin St., Suite 2300  
Houston, Texas 77002-6760  
512.542.8569  
Date: April 21, 2006  
657631\_1.DOC